

Supplement1. Satisfaction with ongoing contraceptive use by BMI (four BMI groups)

	Very Satisfied	Satisfied	Neutral	Dissatisfied	Very dissatisfied	P-value
BMI						
15-18.49 (n=26)	15 (57.7)	4 (15.4)	5 (19.2)	2 (7.7)	0 (0.0)	0.015
18.50-24.99 (n=742)	318 (42.9)	274 (36.9)	104 (14.0)	41 (5.5)	5 (0.7)	
25-29.99 (n=249)	130 (52.2)	72 (28.9)	33 (13.3)	11 (4.4)	3 (1.2)	
30 (n=111)	56 (50.5)	26 (23.4)	18 (16.2)	7 (6.3)	4 (3.6)	

BMI, body mass index.

All data are n (%). P-value calculated with Chi-square test with p-value ≥ 0.05 as significant.

Supplement 2. Satisfaction with ongoing contraceptive use by age (four age groups)

	Very Satisfied	Satisfied	Neutral	Dissatisfied	Very dissatisfied	P-value
Age						
20-24 (n=499)	222 (44.5)	178 (35.7)	71 (14.2)	25 (5.0)	3 (0.6)	0.248
25-29 (n=294)	138 (46.9)	102 (34.7)	35 (11.9)	13 (4.4)	6 (2.0)	
30-34 (n=190)	81 (42.6)	62 (32.6)	31 (16.3)	12 (6.3)	4 (2.1)	
35-40 (n=156)	78 (50.0)	45 (28.8)	20 (12.8)	13 (8.3)	0 (0.0)	

All data are n (%). P-value calculated with Chi-square test with p-value ≥ 0.05 as significant.